



TITLE:

Bleaching of leaf litter and associated microfungi in subboreal and subalpine forests.

AUTHOR(S):

Hagiwara, Yusuke; Matsuoka, Shunsuke; Hobara, Satoru; Mori, Akira S; Hirose, Dai; Osono, Takashi

CITATION:

Hagiwara, Yusuke ...[et al]. Bleaching of leaf litter and associated microfungi in subboreal and subalpine forests.. Canadian journal of microbiology 2015, 61(10): 735-743

ISSUE DATE:

2015-10

URL:

<http://hdl.handle.net/2433/202572>

RIGHT:

This is the accepted manuscript (post-print) of an article is available at <http://dx.doi.org/10.1139/cjm-2015-0111>; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 ; This is not the published version. Please cite only the published version.

- 1 Bleaching of leaf litter and associated microfungi in subboreal and subalpine
- 2 forests
- 3
- 4 Yusuke Hagiwara, Shunsuke Matsuoka, Satoru Hobara, Akira S. Mori, Dai
- 5 Hirose, and Takashi Osono
- 6
- 7 **Y. Hagiwara, S. Matsuoka, T. Osono.** Center for Ecological Research, Kyoto
- 8 University, Otsu, Shiga 520-2113, Japan
- 9 **S. Hobara.** Department of Environmental Symbiotic Sciences, Rakuno Gakuen
- 10 University, Ebetsu, Hokkaido 069-8501, Japan
- 11 **A. S. Mori.** Graduate School of Environment and Information Sciences,
- 12 Yokohama National University, Yokohama, Kanagawa 204-8501, Japan
- 13 **D. Hirose.** College of Pharmacy, Nihon University, Funabashi, Chiba 274-8555,
- 14 Japan
- 15
- 16 **Corresponding author:** Takashi Osono (e-mail: tosono@ecology.kyoto-u.ac.jp)

17

18 **Abstract:** Fungal decomposition of lignin leads to the whitening, or bleaching, of
19 leaf litter, especially in temperate and tropical forests, but less is known about
20 such bleaching in forests of cooler regions, such as boreal and subalpine forests.
21 The purposes of the present study were to examine the extent of bleached area
22 on the surface of leaf litter and its variation with environmental conditions in
23 subboreal and subalpine forests in Japan and microfungi associated with the
24 bleaching of leaf litter by isolating fungi from the bleached portions of the litter.
25 Bleached area accounted for 21.7% to 32.7% and 2.0% to 10.0% of total leaf area
26 of *Q. crispula* and *B. ermanii*, respectively, in subboreal forests, and for 6.3% and
27 18.6% of total leaf area of *B. ermanii* and *P. jezoensis* var. *hondoensis*,
28 respectively, in a subalpine forest. In subboreal forests, elevation, C/N ratio and
29 pH of FH layer, and slope aspect were selected as predictor variables for the
30 bleached leaf area. Leaf mass per area and lignin content were consistently
31 lower in the bleached area than in the nonbleached area of the same leaves,
32 indicating that the selective decomposition of acid unhydrolyzable residue (AUR,

recalcitrant compounds such as lignin, tannins, and cutins) enhanced the mass loss of leaf tissues in the bleached portions. Isolates of a total of 11 fungal species (six species of Ascomycota and five of Basidiomycota) exhibited leaf-litter-bleaching activity under pure culture conditions. Two fungal species (*Coccoomyces* sp. and *Mycena* sp.) occurred in both subboreal and subalpine forests which were separated from each other by approximately 1100 km.

Key words: decomposition, elevational gradient, fungal diversity, leaves, lignin, ligninolytic fungi.

Introduction

Fungi play central roles in the decomposition of lignin and other recalcitrant compounds (often registered as acid unhydrolyzable residues) in leaf litter of forest trees (van der Wal et al. 2013). Fungal decomposition of lignin is mediated by the activity of such extracellular enzymes as lignin peroxidases,

manganese peroxidases, phenol oxidases, and laccases (Eriksson et al. 1990) and often leads to the whitening, or bleaching, of leaf litter (Osono 2007). The fungal bleaching of leaf litter has been reported mainly from temperate and tropical forests (Osono 2006; Osono et al. 2008a, 2009), and less is known about it in forests of cooler regions, such as boreal and subalpine forests. The report of Hintikka (1970) is one of the milestone papers about the diversity and functioning of macrofungi (mainly in the Basidiomycota) associated with the bleaching of forest litter in Finnish boreal forests. More recently, Miyamoto et al. (2000) and Osono (2015c) showed that a suite of macrofungi in the Basidiomycota were capable of actively decomposing lignin and bleaching leaf litter in subboreal and subalpine forests of Japan. Compared to information about macrofungi, however, information is still lacking in boreal and subalpine forests regarding microfungi associated with the bleaching of leaf litter (Osono 2011) and possible climatic and environmental factors affecting their abundance.

The purposes of the present study were to examine (i) the extent of

bleached area on the surface of leaf litter and its variation with environmental conditions and geographic locations and (ii) microfungi associated with the bleaching of leaf litter by isolating fungi from the bleached portions as well as surrounding nonbleached portions of the litter. We selected two study sites, one in subboreal forest in northern Japan and another in subalpine forest in central Japan, separated by 1100 km from each other, but having similar climatic conditions and vegetation. Special attention was paid to an elevational gradient on a mountain slope in the subboreal forest, in which sampling was conducted at five elevational classes from 200 m to 1000 m. Several environmental factors were measured at sampling sites of different elevational classes to test the relative importance of these factors affecting bleaching on leaf litter. Fungi were isolated from bleached areas, as well as nonbleached areas, and were then tested for their bleaching activity under a pure culture condition, and subjected to DNA base sequence analysis for taxonomic identity and assignment.

Materials and Methods

81

82 **Study sites**

83 Samples were collected from two sites in Japan: subboreal forests in
84 Hokkaido and a subalpine forest in Gifu. The subboreal forests were located on
85 the northwest slope of Mt. Rausu (44°04'N, 145°07'E) in Shiretoko Peninsula,
86 northeastern Japan. Mean annual temperature is 6.2 °C, and mean annual
87 precipitation is 1090 mm at Utoro Station of Automatic Meteorological Data
88 Acquisition System of Japan Meteorological Agency, located 3 m above sea level
89 and 10 km southwest of the study site. Dominant trees in the plots included
90 *Abies sachaliensis* F.Schmidt, *Quercus crispula* Bl., *Betula platyphylla* Suk.,
91 and *B. ermanii* Cham. (Mori et al. in press). Bleaching was noticeable on leaf
92 litter of such tree species as *Q. crispula*, *B. ermanii*, *Kalopanax pictus* (Thunb.)
93 Nakai, *Tilia maximowicziana* Shirasawa, *Acer japonicum* Thunb., and *A.*
94 *sachaliensis*. In 2010, 10 study plots (10 m × 10 m) each were established at 200,
95 400, 600, 800, and 1000 m above sea level, making a total of 50 plots (Mori et al.
96 in press), and used for the collection of leaf litter. The maximum distance

97 between the plots at each elevational class ranged between 311 and 435 m.

98 Further details of the study site were described in Ikeda et al. (2014).

99 The subalpine forest was located on the north slope of Mt. Ontake,

100 central Japan (35°56'N, 137°28'E, 2050 m above sea level). Mean annual

101 temperature is approximately 2 °C, and mean annual precipitation is

102 approximately 2500 mm (Osono 2015b). The study site contains mainly four

103 coniferous species [*Abies mariesii* Masters, *A. veitchii* Lindley, *Picea jezoensis*

104 var. *hondoensis* (Sieb. et Zucc.) Carrière, and *Tsuga diversifolia* (Maxim.)

105 Masters] and one hardwood species (*B. ermanii*). Bleaching was noticeable on

106 leaf litter of two tree species: *B. ermanii* and *P. jezoensis* var. *hondoensis*. A

107 study plot of 50 m × 10 m was laid out in 2008 and used for the collection of leaf

108 litter. Further details of the study site were given in Osono and Takeda (2007).

109

110 Collection of leaves

111 Sampling was conducted in the subboreal forests in July 2010. Leaves

112 of *Q. crispula* and *B. ermanii* of which more than half of the original leaf area

113 remained were collected from the surface of the forest floor beneath the canopy
114 of the respective tree species using a 15 cm × 15 cm quadrat. Leaves of *Q.*
115 *crispula* were present in 10 plots each of elevational classes 200, 400, and 600 m,
116 but not in the plots of elevational class 800 or 1000 m. Leaves of *B. ermanii* were
117 collected from 7, 4, 10, 10, and 10 plots of elevational classes 200, 400, 600, 800,
118 and 1000 m, respectively. Additionally, 20 leaves each of *Q. crispula* and *B.*
119 *ermanii* with evident bleaching were arbitrarily collected from the plots at 400
120 m and were used for fungal isolation.

121 Sampling of leaves of *B. ermanii* and *P. jezoensis* var. *hondoensis* was
122 conducted in the subalpine forest in June and August 2008, respectively. Leaves
123 of *B. ermanii* of which more than half of the original leaf area remained, or
124 needles of *P. jezoensis* var. *hondoensis* were collected from the surface of the
125 forest floor beneath the canopy of 10 individuals of each tree species using a 15
126 cm × 15 cm quadrat. Additionally, 20 leaves each of *B. ermanii* and *P. jezoensis*
127 var. *hondoensis* with evident bleaching were arbitrarily collected from the plot
128 and were used for fungal isolation. Trees are referred to by their genus names in

129 the present study for the sake of simplicity.

130

131 **Measurement of leaves**

132 The leaves for leaf area measurement were placed in paper bags and
133 taken back to the laboratory at ambient temperature. The leaves of *Quercus* and
134 *Betula* were pressed between layers of plywood and paper and oven-dried at
135 40 °C for one week. The leaves were then photocopied and scanned with a
136 photocopier (EPSON GT-8000). Total leaf area and the proportion of bleached
137 area were measured with an image software (NIH image, Windows version
138 v.4.0.3, Scion), according to the method described in Hagiwara et al. (2012). Ten
139 *Picea* needles were randomly selected for each quadrat, oven-dried at 40 °C for
140 one week, and measured for total length and the length of needle portions with
141 bleaching under a binocular microscope with magnification of 40× (Hirose and
142 Osono 2006). The bleached area or bleached length of leaves was defined as the
143 bleached percentage of the total leaf area or length. The bleached length of *Picea*
144 needles was denoted here as the bleached area for the sake of simplicity. The

mean value of the bleached area was calculated for each elevational class in subboreal forests and for the plot in the subalpine forest.

A 6-mm-diameter cork borer was used to excise leaf disks, avoiding the primary vein, from the bleached area and surrounding nonbleached area of the same leaves of *Q. crispula* and *B. ermanii* from subboreal and subalpine forests. Three to 61 disks (mean: 11 disks) were punched out for each quadrat. The disks were oven-dried again at 40 °C for one week and weighed to calculate leaf mass per area (LMA). To compare the chemical composition, the disks were combined to make one sample each of bleached and nonbleached leaf area for each tree species and each study site, ground in a laboratory mill, and used for proximate chemical analyses. Content of acid unhydrolyzable residue (AUR) was measured with hot sulfuric acid digestion, according to the method described in Osono (2015c).

We measured five environmental variables for each study plot of subboreal forests that were expected to influence the occurrence of bleached portions on leaves: carbon to nitrogen (C/N) ratio, pH (KCl), and gravimetric

water content of all materials in FH layer, inclination of the slope, and the slope aspect. Methods of measurement of these variables were described in Mori et al. (in press). Briefly, total carbon and nitrogen contents were determined with combustion method by automatic gas chromatography (NC analyzer SUMIGRAPH NC-900, Sumitomo Chemical, Osaka, Japan). The pH was measured in a 1 N potassium chloride solution with Docu-pH meter+ (Sartorius, Goettingen, Germany).

Isolation and screening of bleaching fungi

The leaves for fungal isolation were placed in paper bags and preserved at 4 °C for no longer than three days before the isolation of fungi. Single leaf disks were excised from the bleached portion of individual leaves of *Quercus* and *Betula*, using a sterile 6-mm-diameter cork borer, and other single disks were excised from the nonbleached portion of the same leaves. One bleached and one nonbleached portion (5-10 mm in length) were cut separately and aseptically for individual *Picea* needles. This produced a total of 20 bleached and 20

177 nonbleached disks or needle portions for each tree species from each study site.

178 Fungi were isolated from bleached and nonbleached disks or needle portions

179 using the surface disinfection method according to Hirose et al. (2014). The

180 surface-disinfected materials were plated on 9-cm Petri dishes containing 2%

181 lignocellulose agar (LCA) modified as described by Miura and Kudo (1970)

182 [glucose 0.1%, KH_2PO_4 0.1%, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02%, KCl 0.02%, NaNO_3 0.2%,

183 yeast extract 0.02%, and agar 2% (w/v)], two disks or needle portions per plate.

184 Note that the modified LCA of Miura and Kudo (1970) does not contain lignin or

185 other recalcitrant compounds. The modified LCA was used because its low

186 glucose content suppresses the overgrowth of fast-growing fungal species (Osono

187 and Takeda 1999). The plates were incubated in darkness at 10 °C and observed

188 for 4 weeks after the disinfection. Any fungal hyphae or spores appearing on the

189 plates were subcultured onto fresh LCA plates, incubated, and observed

190 micromorphologically.

191 Isolates were tested for their ability to bleach leaf litter under a pure

192 culture condition. Newly shed leaves of *Quercus* and *Betula* without obvious

193 fungal or faunal attack were collected from the forest floor of subboreal forests
194 in July 2010, and those of *Betula* were collected from the subalpine forest in
195 September 2008. These leaves were oven-dried at 40°C for one week, cut into
196 pieces (1 cm × 1 cm), and autoclaved at 120°C for 20 min. The sterilized leaf
197 pieces were placed on the surface of Petri dishes (6 cm diameter), one piece per
198 plate, containing 20 ml of 2% malt extract agar [malt extract 2% and agar 2%
199 (w/v)] previously inoculated with fungal isolates and incubated at 20°C for two
200 weeks. *Quercus* isolates were inoculated with sterilized *Quercus* leaf pieces, and
201 *Betula* and *Picea* isolates were inoculated with *Betula* leaf pieces. The plates
202 were sealed firmly with laboratory film so that moisture did not limit the
203 bleaching on the agar medium and incubated at 20°C for 24 weeks in the dark.
204 After incubation, the leaf pieces were retrieved and examined under a binocular
205 microscope with magnification of 40× for the occurrence of bleaching on the leaf
206 surface. The occurrence of bleaching was scored visually into six classes: 0, no
207 bleached portion; 1, bleached in 1% to 20% of leaf area; 2, bleached in 21% to
208 40%; 3, bleached in 41% to 60%; 4, bleached in 61% to 80%; and 5, bleached in

209 81% to 100%. The visual class was transformed into the percentage bleached area
210 with respect to the total leaf area by applying the median for each of class 1 to 5.
211 Three control plates with the sterilized leaves but no fungal inoculation were
212 also incubated under the same conditions. No microbial colonies or bleaching
213 developed on the control plates, indicating the effectiveness of the sterilization
214 method used in the present study. Fungal isolates with the score of 1 to 5 were
215 regarded as possessing bleaching activity. We obtained a total of 180 fungal
216 isolates from 160 leaf disks (four litter types \times two portions \times 20 leaves), of
217 which 40 isolates were judged to be bleaching fungi and used for molecular
218 analysis as described below.

219

220 **Molecular methods**

221 Genomic DNA was extracted from mycelia that had been cultured on
222 2.5% malt extract agar overlaid with a cellophane membrane following the
223 modified CTAB method described by Matsuda and Hijii (1999). Polymerase
224 chain reaction (PCR) was performed using a Quick Taq HS DyeMix (Toyobo,

225 Osaka, Japan). Each PCR reaction contained a 50 μ l mixture [21 μ l distilled
226 water, 25 μ l master mix, 3 μ l ca. 0.5 ng/ μ l template DNA, and 0.5 μ l of each
227 primer (final, 0.25 μ M)]. To PCR amplify the region including the rDNA ITS and
228 28S rDNA D1-D2 domain, the primer pair ITS1f (Gardes and Bruns 1993) and
229 LR3 (Vilgalys and Hester 1990) was used. Each DNA fragment was amplified
230 using a PCR thermal cycler (DNA engine; Bio-Rad, Hercules, CA, USA) using
231 the following thermal cycling schedule. The first cycle consisted of 5 min at 94°C,
232 followed by 35 cycles of 30 s at 94°C, 30 s at 50°C for annealing, 1 min at 72°C,
233 and a final cycle of 10 min at 72°C. The reaction mixture was then cooled at 4°C
234 for 5 min. PCR products were purified with a QIAquick PCR Purification Kit
235 (Qiagen, Germany) according to the manufacturer's instructions.

236 Purified PCR products were sequenced by FASMAC Co., Ltd.
237 (Kanagawa, Japan). Sequencing reactions were performed in a Gene Amp PCR
238 System 9700 (Applied Biosystems, USA) using a BigDye Terminator V3.1
239 (Applied Biosystems), following the protocols supplied by the manufacturer. The
240 fluorescent-labeled fragments were purified from the unincorporated

241 terminators using an ethanol precipitation protocol. The samples were
242 resuspended in formamide and subjected to electrophoresis in an ABI 3730xl
243 sequencer (Applied Biosystems).

244 The sequences determined in the present study were deposited in the
245 DNA Data Bank of Japan (DDBJ) (LC014885 to LC014897). The rDNA ITS and
246 28S sequences were compared with available rDNA sequences in the GenBank
247 database using BLASTN searches (Altschul et al. 1990). For phylogenetic
248 analysis, MAFFT ver. 7 (Kato and Standley 2013) was used for preliminary
249 multiple alignments of nucleotide sequences. Final alignments were manually
250 adjusted using BioEdit (Hall 1999). Alignment gaps were treated as missing
251 data, and ambiguous positions were excluded from the analysis. Phylogenetic
252 tree was conducted by Maximum Likelihood (ML) methods (Felsenstein 1981)
253 with the best fit nucleotide substitution model based on the lowest BIC score
254 (Bayesian Information Criterion). To estimate clade support, the bootstrap
255 procedure of Felsenstein (1985) was employed with 1000 replicates. These
256 analyses were carried out using MEGA6 (Tamura et al. 2013).

257 Based on the phylogenetic tree, the isolates were manually grouped into
258 operational taxonomic units according to the similarity of 28S rDNA D1-D2
259 sequences with a criterion of 98%. Based on the results of molecular barcoding
260 and micromorphological observation, we determined the identity and taxon of
261 fungal isolates with bleaching activity.

262

263 **Statistical analysis**

264 Generalized linear models (GLMs) were used to evaluate the difference
265 in bleached area of *B. ermanii* and *Q. crispula* leaf litter in subboreal forests
266 with a Gaussian distribution using the elevational class, C/N ratio, pH, and
267 water content of FH layer, the inclination of the slope, and the slope aspect as
268 predictor variables. The GLMs were performed with the *glm* function of R
269 version 3.0.2 for Mac (<http://www.r-project.org>) and with the *glht* function of the
270 R multcomp package for multiple comparisons with Tukey's test. To find the
271 most parsimonious models we performed automatic forward stepwise model
272 selections with the Akaike information criterion (AIC), using the minimum AIC

as the best-fit estimator. The model selections were performed with the stepAIC function of R. The GLMs were used to evaluate the difference in LMA of *B. ermanii* and *Q. crispula* leaf litter in subboreal forests with a Gaussian distribution using the leaf portions (bleached vs nonbleached) and the elevational class as predictor variables.

Results

Bleached area on leaf litter

Bleached area accounted for 21.7% to 32.7% and 2.0% to 10.0% of the mean total leaf area of *Q. crispula* and *B. ermanii*, respectively, in subboreal forests and for 6.3% and 18.6% of total leaf area of *B. ermanii* and *P. jezoensis* var. *hondoensis*, respectively, in subalpine forest (Fig. 1a). In subboreal forests, elevational class, C/N ratio and pH of FH layer, and/or slope aspect were selected as predictor variables for the bleached leaf area of *Q. crispula* and *B. ermanii*, of which the elevational class and C/N ratio of FH layer had significant

289 χ^2 values ($P < 0.05$) for *Q. crispula* and the elevational class had a significant χ^2
290 value ($P < 0.05$) for *B. ermanii* (Table 1, Fig. 1b). The bleached area on *Q. crispula*
291 leaves was significantly ($P < 0.05$) greater at 600 m than at 200 and 400 m, and
292 that on *B. ermanii* leaves was significantly ($P < 0.05$) greater at 600 m than at
293 800 and 1000 m. The water content of FH layer and the inclination of the slope
294 were not selected as predictor variables for the bleached leaf area of *Q. crispula*
295 or *B. ermanii*.

296 Leaf mass per area (LMA) was consistently lower in the bleached area
297 than in the nonbleached area of the same leaves for *Q. crispula* and *B. ermanii*
298 from subboreal and subalpine forests (Table 2). The difference in LMA between
299 the bleached and nonbleached area was statistically significant (GLM, $P < 0.001$)
300 for the two tree species from subboreal forests (Table 2). The LMA of *Q. crispula*
301 was also significantly (GLM, $P < 0.001$) lower at the elevational classes of 400
302 and 600 m than at 200 m (Table 2). AUR content was lower in the bleached leaf
303 area than in the nonbleached leaf area for *Q. crispula* and *B. ermanii* from
304 subboreal forests (Table 2).

305

306 Fungi with bleaching activity

307 A total of 11 fungal species (six in Ascomycota and five in
308 Basidiomycota) were isolated from bleached and/or nonbleached portions and
309 exhibited bleaching activity under pure culture conditions (Table 3, Fig. 2). Five
310 of the six ascomycete species belonged to Rhytismataceae and the other to
311 Xylariaceae; two basidiomycete species belonged to Tricholomataceae and the
312 others to Mycenaceae, Russulaceae, or Phanerochaetaceae (Fig. 2). The
313 bleaching activity was lower in Xylariaceae sp. (which caused mean bleaching of
314 less than 10% of the leaf area) than in rhytismataceous and basidiomycete
315 species (which causing bleaching of more than or equal to 50% of the leaf area)
316 (Table 3).

317 Three, three, five, and four fungal species were detected as fungi with
318 bleaching activity toward *Q. crispula*, *B. ermanii* of subboreal forest, *B. ermanii*
319 of subalpine forest, and *P. jezoensis* var. *hondoensis*, respectively (Table 3).
320 *Coccoomyces* sp. and *Mycena* sp. occurred on leaves of three tree species in both

321 subboreal and subalpine forests (Table 3).

322

323 Discussion

324

325 The proportions of the bleached area of *Q. crispula* leaves were at the
326 highest part of the range previously reported for tree leaves in temperate and
327 tropical forests, whereas those of *B. ermanii* leaves were at the low to middle
328 part of the range (summarized in Osono 2006; Hagiwara et al. 2012). The higher
329 bleached area of *Q. crispula* leaves than in that of *B. ermanii* leaves was possibly
330 attributed to the lower content of AUR (Table 2) and nitrogen (discussed further
331 below) in *Q. crispula* leaves. Few data have been available on the bleached
332 portion of conifer needles comparable to the needles of *P. jezoensis* var.
333 *hondoensis* in the subalpine forest. The lower LMA and the lower AUR content
334 on the bleached area than on the nonbleached area are consistent with previous
335 data (Osono 2006) and indicate that the selective decomposition of AUR
336 enhanced the mass loss of leaf tissues in the bleached portions which

337 ligninolytic fungi colonized, compared to the nonbleached portions of the same
338 leaves.

339 In subboreal forests, the proportions of the bleached area on leaves of
340 both *Q. crispula* and *B. ermanii* reached the highest value at the elevational
341 class of 600 m, and that of *Q. crispula* was significantly affected by the C/N ratio
342 of FH layer (Fig. 1, Table 1). These results suggested that the bleached area did
343 not respond linearly with the temperature lapse with the elevation and possible
344 effects of other local factors. For example, previous studies demonstrated
345 difference in microfungal assemblages at different elevational classes (van
346 Maanen et al. 2000; Gourbière et al. 2001; Osono and Hirose 2009b). There is
347 also some evidence to suggest that high N content could inhibit colonization by
348 ligninolytic fungi. For example, N could cause a biochemical suppression of
349 AUR-degrading enzymes of fungi (Keyser et al. 1978; Fenn et al. 1981). Osono et
350 al. (2002) reported that avian excreta rich in N suppressed the colonization of
351 litter by ligninolytic fungi. Similarly, Hagiwara et al. (2012) found that the
352 bleached area of *Acacia mangium* leaf litter was negatively correlated with N

content of the litter, supporting the notion that the colonization of leaves by ligninolytic fungi could be inhibited by higher N content. The unimodal pattern of bleached area on *B. ermanii* leaves along the elevational gradient (Fig. 1a) implied that not only N level, pH, and the slope aspect but also other microenvironmental factors, such as the composition of tree and ground vegetation, singly or in combination, might affect the occurrence of bleached area on the litter. Two other possible factors affecting the occurrence of bleached portions on leaf litter along the elevational gradient include (i) different leaf fall time between the study sites and (ii) competition between bleaching fungi and other microbes.

The present results of proximate chemical composition analysis of leaf tissues, the isolation of fungi, and the pure culture bleaching assay demonstrated that 11 fungal species were responsible for AUR decomposition and bleaching of leaf litter in the subboreal and subalpine forests. The taxonomic composition of bleaching fungi of subboreal and subalpine leaves was generally consistent with that reported for other temperate and tropical regions.

369 Regarding Ascomycetes, species in the Rhytismataceae have been reported to
370 bleach the leaf litter and decompose the AUR fraction of *Camellia japonica*
371 (Koide et al. 2005b; Osono and Hirose 2009a), *Gaultheria shallon* (Osono et al.
372 2008b), and *Pinus* spp. (Osono and Hirose 2011). These rhytismataceous species
373 can be endophytic on live leaf tissues and take part in lignin decomposition in
374 early stages of decomposition (Koide et al. 2005a). Fungi in the Xylariaceae in
375 Ascomycetes are well known for their ligninolytic activity, especially in the
376 tropics (e.g. Osono et al. 2011, 2013). Basidiomycetes are major components of
377 ligninolytic fungal assemblages in temperate and tropical forests (Boddy et al.
378 2008; Osono 2015a). *Mycena* sp. isolated as a bleaching fungus in the present
379 study was also found fruited on the forest floor of the subalpine forest (Osono
380 2015a) and caused the selective delignification of *Betula* leaves (Osono 2015d).

381 The present study demonstrated the occurrence of bleaching on leaf
382 litter of subboreal and subalpine forests in Japan, and the association of fungi
383 with it. The occurrence of the same fungal species (*Coccomyces* sp. and *Mycena*
384 sp.) at the two locations, despite their distance of approximately 1100 km from

each other, can be partly attributed to the similarity in the climatic conditions and vegetation between these subboreal and subalpine forests. Similarly, fruiting bodies of such ligninolytic macrofungi as *M. aurantiidisca* were observed in both a subboreal forest in Hokkaido (Miyamoto et al. 1998) and the subalpine forest of the present study (Osono 2015a). It remains unclear to what extent the bleaching fungi contribute to the loss of AUR in decomposing litter on the forest floor of these forest stands. Tian et al. (2000), for example, found that the decomposition of AUR was slower than that of carbohydrates in leaf litter during a 3.5-year incubation in subalpine forest, suggesting relatively minor roles of ligninolytic activity compared to cellulolytic activity of the whole fungal assemblages. Further studies will be needed to follow simultaneously the changes in bleached area, the succession of ligninolytic fungi, and the loss of AUR in decomposition processes to quantify the role of fungi associated with the bleaching.

Acknowledgments

401
402 This study was supported by the Ministry of Education, Culture, Sports, Science,
403 and Technology of Japan (MEXT) (No.19780114, No.15K07480), the Sumitomo
404 Foundation, and the Japan Securities Scholarship Foundation. The Shiretoko
405 Foundation provided logistical support for the fieldwork. We thank H. Kondo, K.
406 Koizumi, Y. Miyake, W. Azuma, H. Kamauchi, and N. Mori for their assistance.

407

408 **References**

409

- 410 Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. 1990. Basic
411 local alignment search tool. *J. Mol. Biol.* **215**: 403-410.
- 412 Boddy, L., Frankland, J.C., and van West, P. 2008. Ecology of Saprotrophic
413 Basidiomycetes. Academic Press, London.
- 414 Eriksson, K.E., Blanchetter, R.A., and Ander, P. 1990. Microbial and enzymatic
415 degradation of wood and wood components. Springer, New York.
- 416 Felsenstein, J. 1981. Evolutionary trees from DNA sequences: a maximum

- 417 likelihood approach. *J. Mol. Evol.* **17**: 368–376.
- 418 Felsenstein, J, 1985. Confidence limits on phylogenies: an approach using the
- 419 bootstrap. *Evolution* **39**: 783–791.
- 420 Fenn, P., Choi, S., and Kirk, T.K. 1981. Ligninolytic activity of *Phanerochaete*
- 421 *chrysosporium*: Physiology of suppression by NH_4^+ and L-Glutamate.
- 422 *Microbiology* **130**: 66-71.
- 423 Gardes, M., and Bruns, T.D. 1993. ITS primers with enhanced specificity for
- 424 basidiomycetes: application to the identification of mycorrhizae and
- 425 rusts. *Mol. Ecol.* **21**: 113-118.
- 426 Gourbière, F., van Maanen, A., and Debouzie, D. 2001. Associations between
- 427 three fungi on pine needles and their variation along a climatic gradient.
- 428 *Mycol. Res.* **105**: 1101-1109.
- 429 Hagiwara, Y., Osono, T., Ohta, S., Agus, W., and Hardjono, A. 2012. Colonization
- 430 and decomposition of leaf litter by ligninolytic fungi in *Acacia mangium*
- 431 plantations and adjacent secondary forests. *J. For. Res.* **17**: 51-57.
- 432 Hall, TA, 1999. BioEdit: a user-friendly biological sequence alignment editor and

- 433 analysis program for Windows 95/98/NT. Nucleic Acids Symp. Ser. **41**:
- 434 95-98.
- 435 Hintikka, V. 1970. Studies on white-rot humus formed by higher fungi in forest
- 436 soils. Commun. Inst. For. Fenn. **69.2**: 1-68.
- 437 Hirose, D., and Osono, T. 2006. Development and seasonal variation of
- 438 *Lophodermium* populations on *Pinus thunbergii* needle litter.
- 439 Mycoscience 47: 242-247.
- 440 Hirose, D., Tanabe, Y., Uchida, M., Kudoh, S., and Osono, T. 2014. Microfungi
- 441 associated with withering willow wood in ground contact near Syowa
- 442 Station, East Antarctica for 40 years. Polar Biol. **36**: 919-924.
- 443 Ikeda, A., Matsuoka, S., Masuya, H., Mori, A.S., Hirose, D., and Osono, T. 2014.
- 444 Comparison of the diversity, composition, and host recurrence of
- 445 xylariaceous endophytes in subtropical, cool temperate, and subboreal
- 446 regions in Japan. Popul. Ecol. **56**: 289-300.
- 447 Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software
- 448 version 7: improvements in performance and usability. Mol. Biol. Evol.

- 449 30: 772–780.
- 450 Keyser, P., Kirk, T.K., and Zeikus, J.G. 1978. Ligninolytic enzyme system of
- 451 *Phanerochaete chrysosporium*: Synthesized in the absence of lignin in
- 452 response to nitrogen starvation. J. Bacteriol. **135**: 790-797.
- 453 Kimura, M., 1980. A simple method for estimating evolutionary rate of base
- 454 substitutions through comparative studies of nucleotide sequences. J.
- 455 Mol. Evol. **16**: 111-120.
- 456 Koide, K., Osono, T., and Takeda, H. 2005a. Colonization and lignin
- 457 decomposition of *Camellia japonica* leaf litter by endophytic fungi.
- 458 Mycoscience **46**: 280-286.
- 459 Koide, K., Osono, T., and Takeda, H. 2005b. Fungal succession and
- 460 decomposition of *Camellia japonica* leaf litter. Ecol. Res. **20**: 599-609.
- 461 Matsuda, Y., and Hijii, N. 1999. Characterization and identification of
- 462 Strobilomyces confusus ectomycorrhizas on momi fir by RFLP analysis
- 463 of the PCR-amplified ITS region of the rDNA. J. For. Res. **4**: 145-150.
- 464 Miyamoto, T., Igarashi, T., and Takahashi, K. 1998. Notes on three species of

- 465 *Mycena* new to Japan from *Picea* forests of Hokkaido. *Mycoscience* **39**:
- 466 337-342.
- 467 Miyamoto, T., Igarashi, T., and Takahashi, K. 2000. Lignin-degrading ability of
- 468 litter-decomposing basidiomycetes from *Picea* forests of Hokkaido.
- 469 *Mycoscience* 41: 105-110.
- 470 Miura, K., and Kudo, M. 1970. An agar-medium for aquatic hyphomycetes.
- 471 *Trans. Mycol. Soc. Japan* **11**: 116-118. (in Japanese)
- 472 Mori, A.S., Shiono, T., Haraguchi, T., Ota, A., Koide, D., Ohgue, T., Kitagawa, R.,
- 473 Maeshiro, R., Aung, T.T., Nakamori, T., Hagiwara, Y., Matsuoka, S.,
- 474 Ikeda, A., Hishi, T., Hobara, S., Mizumachi, E., Frisch, A., Thor, G., Fujii,
- 475 S., Osono, T., and Gustafsson, L. Functional redundancy of multiple
- 476 forest taxa along an elevational gradient: predicting the consequences of
- 477 non-random species loss. *J. Biogeogr.* in press. doi: 10.1111/jbi.12514
- 478 Osono, T. 2006. Fungal decomposition of lignin in leaf litter: comparison
- 479 between tropical and temperate forests. *In* Proceedings for the 8th
- 480 International Mycological Congress, August 20-25, 2006, Cairns,

- 481 Australia. *Edited by* W. Meyer and C. Pearce. Medimond, Italy. pp.
- 482 111-117.
- 483 Osono, T. 2007. Ecology of ligninolytic fungi associated with leaf litter
- 484 decomposition. *Ecol. Res.* **22**: 955-974.
- 485 Osono, T. 2011. Diversity and functioning of fungi associated with leaf litter
- 486 decomposition in an Asian climatic gradient. *Fungal Ecol.* **4**: 375-385.
- 487 Osono, T. 2015a. Diversity, resource utilization, and phenology of fruiting bodies
- 488 of litter-decomposing macrofungi in subtropical, temperate, and
- 489 subalpine forests. *J. For. Res.* **20**: 60-68.
- 490 Osono, T. 2015b. Hyphal length in the forest floor and soil of subtropical,
- 491 temperate, and subalpine forests. *J. For. Res.* **20**: 69-76.
- 492 Osono, T. 2015c. Effects of litter type, origin of isolate, and temperature on
- 493 decomposition of leaf litter by macrofungi. *J. For. Res.* **20**: 77-84.
- 494 Osono, T. 2015d. Decomposing ability of diverse litter-decomposer macrofungi in
- 495 subtropical, temperate, and subalpine forests. *J. For. Res.* **20**: in press.
- 496 Osono, T., and Takeda, H. 1999. A methodological survey on incubation of fungi

- 497 on leaf litter of *Fagus crenata*. Ap. For. Sci. Kansai **8**: 103-108. (in
498 Japanese with English abstract)
- 499 Osono, T., and Takeda, H. 2007. Microfungi associated with *Abies* needles and
500 *Betula* leaf litter in a subalpine coniferous forest. Can. J. Microbiol. **53**:
501 1-7.
- 502 Osono, T., and Hirose, D. 2009a. Effects of prior decomposition of *Camellia*
503 *japonica* leaf litter by an endophytic fungus on the subsequent
504 decomposition by fungal colonizers. Mycoscience **50**: 52-55.
- 505 Osono, T., and Hirose, D. 2009b. Altitudinal distribution of microfungi
506 associated with *Betula ermanii* leaf litter on Mt. Rishiri, northern Japan.
507 Can. J. Microbiol. **55**: 783-789.
- 508 Osono, T., and Hirose, D. 2011. Colonization and lignin decomposition of pine
509 needle litter by *Lophodermium pinastri*. Forest Pathol. **41**: 156-162.
- 510 Osono, T., Hobara, S., Fujiwara, S., Koba, K., and Kameda, K. 2002. Abundance,
511 diversity, and species composition of fungal communities in a temperate
512 forest affected by excreta of the Great Cormorant *Phalacrocorax carbo*.

- 513 Soil Biol. Biochem. **34**: 1537-1547.
- 514 Osono, T., Ishii, Y., and Hirose, D. 2008a. Fungal colonization and decomposition
- 515 of *Castanopsis sieboldii* leaf litter in a subtropical forest. Ecol. Res. **23**:
- 516 909-917.
- 517 Osono, T., Iwamoto, S., and Trofymow, J.A. 2008b. Colonization and
- 518 decomposition of salal (*Gaultheria shallon*) leaf litter by fungi in
- 519 successional forests on coastal British Columbia. Can. J. Microbiol. **54**:
- 520 427-434.
- 521 Osono, T., Ishii, Y., Takeda, H., Seramethakun, T., Khamyong, S., To-Anun, C.,
- 522 Hirose, D., Tokumasu, S., and Kakishima, M. 2009. Fungal succession
- 523 and lignin decomposition on *Shorea obtusa* leaves in a tropical seasonal
- 524 forest in northern Thailand. Fungal Div. **36**: 101-119.
- 525 Osono, T., To-Anun, C., Hagiwara, Y., and Hirose, D. 2011. Decomposition of
- 526 wood, petiole, and leaf litter by *Xylaria* species from northern Thailand.
- 527 Fungal Ecol. **4**: 210-218.
- 528 Osono, T., Tateno, O., and Masuya, H. 2013. Diversity and ubiquity of

- 529 xylariaceous endophytes in live and dead leaves of temperate forest trees.
- 530 Mycoscience **54**: 54-61. Tamura, K., Stecher, G., Peterson, D., Filipski, A.,
- 531 Kumar, S., 2013. MEGA6: molecular evolutionary genetics analysis
- 532 version 6.0. Mol. Biol. Evol. **30**: 2725–2729.
- 533 Tian, X., Takeda, H., and Azuma, J.I. 2000. Dynamics of organic-chemical
- 534 components in leaf litters during a 3.5-year decomposition. Eur. J. Soil
- 535 Biol. **36**: 81-89.
- 536 Van der Wal, A., Geydan, T.D., Kuyper, T.W., and de Boer, W. 2013. A thready
- 537 affair: linking fungal diversity and community dynamics to terrestrial
- 538 decomposition processes. FEMS Microbiol. Rev. **37**: 477–494.
- 539 Van Maanen, A., Debouzie, D., and Gourbière, F. 2000. Distribution of three
- 540 fungi colonizing fallen *Pinus sylvestris* needles along altitudinal
- 541 transects. Mycol. Res. **104**: 1133-1138.
- 542 Vilgalys, R., and Hester, M. 1990. Rapid genetic identification and mapping of
- 543 enzymatically amplified ribosomal DNA from several *Cryptococcus*
- 544 species. J. Bacteriol. **172**: 4238–4246.

545

546 Figure legend

547

548 Fig. 1. Bleached area (% total leaf area or length) on leaves of subboreal and
549 subalpine forests of different elevations (a) and as related to C/N ratio of FH
550 layer in subboreal forests (b). Values are means (n=10) and bars indicate
551 standard errors. Numbers indicate the elevations. Gray columns and open
552 squares, *Betula ermanii*; blank columns and filled circles, *Quercus crispula*;
553 striped columns, *Picea jezoensis* var. *hondoensis*.

554

555 Fig. 2. Maximum-likelihood (ML) phylogeny inferred from 28S rDNA partial
556 sequences including 41 bleaching fungi isolated from bleached and nonbleached
557 portions of leaf litter in subboreal and subalpine forests. The evolutionary model
558 used was the Kimura 2-parameter model (Kimura 1980) with a discrete Gamma
559 distribution (+G, parameter = 0.5624). Bootstrap values for the ML analysis are
560 indicated for corresponding branches. GenBank accession numbers are given in

561 parentheses.

562

Fig. 1






Fig. 2



1 Hagiwara et al. Table 1.

2

3

4

5 Table 1. Summary of generalized linear models for bleached leaf area in subboreal forests
6 and selected predictor variables. ns, not selected. *** P<0.001, * P<0.05. The water content
7 of FH layer and the inclination of the slope were not selected as predictor variables

	<i>Quercus crispula</i>		<i>Betula ermanii</i>	
	Deviance	Probability	Deviance	Probability
Elevational class	772.0	0.02*	463.7	<0.001***
C/N ratio of FH layer	436.3	0.04*	47.3	0.08
pH of FH layer	120.4	0.28	ns	
Slope aspect	ns		31.1	0.16

8

9 Hagiwara et al. Table 2

10

11

12

13 Table 2. Leaf mass per area (LMA, mg/cm²) and the content (mg/g) of acid unhydrolyzable residue (AUR) in bleached (BL) and
14 nonbleached (NB) portions of leaf litter from subboreal and subalpine forests. Values indicate means \pm standard errors
15 (n=4-10). nd, no data. na, not analyzed because of insufficient amount of samples for analysis.

Subboreal					Subalpine	
<i>Quercus crispula</i>			<i>Betula ermanii</i>		<i>Betula ermanii</i>	
BL	NB		BL	NB	BL	NB
LMA in elevational class						
200 m	5.3 \pm 0.3	7.4 \pm 0.4	3.8 \pm 0.3	5.3 \pm 0.6	nd	nd
400 m	4.4 \pm 0.3	6.4 \pm 0.3	3.9 \pm 0.2	5.5 \pm 0.2	nd	nd
600 m	4.1 \pm 0.3	6.1 \pm 0.3	4.4 \pm 0.3	6.3 \pm 0.8	nd	nd
800 m	nd	nd	4.5 \pm 0.6	5.6 \pm 0.8	nd	nd
1000 m	nd	nd	4.4 \pm 0.4	7.3 \pm 0.7	nd	nd
2050 m	nd	nd	nd	nd	4.4 \pm 0.1	5.3 \pm 0.1
Chemical content						
AUR	297.0	442.3	398.2	497.3	na	na

16

17 Table 3. Number of isolates of bleaching fungi isolated from bleached and nonbleached portions of leaf litter and their
18 bleaching activity (as % bleached leaf area caused by fungal isolates, with respect to total leaf area).

	Subboreal		Subalpine		Total number of isolates	Bleaching activity Mean (min-max)	
Fungus	<i>Quercus</i>	<i>Betula</i>	<i>Betula</i>	<i>Picea</i>			
Ascomycota							
<i>Coccomyces</i> sp.	3	1		3	0	7	61 (10-90)
Rhytismataceae sp.1	1	0		0	0	1	90
Xylariaceae sp.	0	6		0	0	6	7 (0-10)
Rhytismataceae sp.2	0	0		3	0	3	63 (50-90)
Rhytismataceae sp.3	0	0		0	10	10	53 (30-90)
<i>Lophodermium</i> sp.	0	0		0	1	1	50
Basidiomycota							
<i>Mycena</i> sp.	0	2		1	4	7	90 (90-90)
<i>Clitocybe</i> sp.	1	0		0	0	1	70
<i>Peniophora</i> sp.	0	0		1	0	1	90
<i>Ceriporia</i> sp.	0	0		1	0	1	50
Tricholomatacea sp.	0	0		0	3	3	90 (90-90)
Number of species	3	3		5	4		